

Hellenic Complex Systems Laboratory

# Relation of Diagnostic Accuracy Measures

Technical Report XIV

Theodora Chatzimichail and Aristides T. Hatjimihail  
2018



## Relation of Diagnostic Accuracy Measures


Theodora Chatzimichail <sup>a</sup> and Aristides T. Hatjimihail <sup>a</sup>

<sup>a</sup> Hellenic Complex Systems Laboratory


**Search Terms:** sensitivity, specificity, diagnostic test, clinical accuracy, diagnostic accuracy, normal distribution, binormal distribution, positive predictive value, negative predictive value, likelihood ratio, odds ratio

### Short Description of the Demonstration

This Demonstration examines the relation of pairs of accuracy measures of diagnostic tests applied on normally distributed nondiseased and diseased populations. This is done for differing prevalence of the disease, taking into account the means and standard deviations of the populations. The means and standard deviations are expressed in arbitrary units. The measures considered are the positive predictive value ("PPV"), the negative predictive value ("NPV"), the (diagnostic) odds ratio ("OR"), the likelihood ratio for a positive result ("LR+") and the likelihood ratio for a negative result ("LR-"). The measures can be selected by clicking the respective "plot" and "versus" buttons.



**Figure 1:** Positive predictive value versus negative predictive value curve of a diagnostic test, with the settings shown at the left.



**Figure 2:** Positive predictive value versus likelihood ratio for a negative result curve of a diagnostic test, with the settings shown at the left.





Figure 3: Odds ratio versus negative predictive value curve of a diagnostic test, with the settings shown at the left.



**Figure 4:** Likelihood ratio for a positive result versus value likelihood ratio for a negative result curve of a diagnostic test, with the settings shown at the left.

## Details

These measures of the clinical accuracy of a diagnostic test applied to a diseased and a nondiseased population can be calculated as functions of the sensitivity and the specificity of the test. Sensitivity is the fraction of the diseased population with a positive test result, while specificity is the fraction of the nondiseased population with a negative test result. If we denote by *sens* the sensitivity, *spec* the specificity, and *pr* the prevalence, then:

$$PPV = \frac{sens \times pr}{spec \times pr + (1 - spec)(1 - pr)}$$

$$NPV = \frac{spec (1 - pr)}{spec (1 - pr) + (1 - sens) pr}$$

$$OR = \frac{\frac{sens}{1 - sens}}{\frac{1 - spec}{spec}}$$

$$LR+ = \frac{sens}{1 - spec}$$

$$LR- = \frac{1 - sens}{spec}$$

Given a diseased and a nondiseased population, the specificity can be defined as a function of sensitivity; therefore, we can generate the parametric plots of any pair of the diagnostic accuracy measures.

To the best of our knowledge, with the exception of the pair PPV and NPV [1, 2], the relation of any other pair of the given diagnostic accuracy measures has not been discussed in the literature.

In the thumbnail, the population data describes a bimodal distribution of serum glucose measurements in nondiabetic and diabetic populations [3].

This Demonstration is an extended version of another Demonstration [2], first published in 2015, and is appropriate as an educational tool for medical students and as an exploratory research tool in diagnostic accuracy studies.

## References

[1] S.-Y. Shiu and C. Gatsonis. The Predictive Receiver Operating Characteristic Curve for the Joint Assessment of the Positive and Negative Predictive Values. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 366(1874), 2008 pp. 2313–2333. doi:10.1098/rsta.2008.0043.

[2] Chatzimichail T. Correlation of Positive and Negative Predictive Values. Wolfram Demonstrations Project, Champaign: Wolfram Research, Inc., 2018 [First published in 2015]. Available at: <http://demonstrations.wolfram.com/CorrelationOfPositiveAndNegativePredictiveValuesOfDiagnostic/>

[3] T.-O. Lim, R. Bakri, Z. Morad and M. A. Hamid. Bimodality in Blood Glucose Distribution: Is It Universal? *Diabetes Care*, 25(12), 2002 pp. 2212–2217. doi:10.2337/diacare.25.12.2212.

## Source Code

Programming language: Wolfram Language

Availability: The updated source code is available at:

<https://www.hcsl.com/Tools/Demonstrations/RelationOfDiagnosticAccuracyMeasures.nb>

## Software Requirements

Operating systems: Microsoft Windows, Linux, Apple iOS

Other software requirements: Wolfram Player®, freely available at: <https://www.wolfram.com/player/> or Wolfram Mathematica®.

## System Requirements

Processor: x86-64 compatible CPU.

System memory (RAM): 4GB+ recommended.

## Permanent Citation:

Chatzimichail T, Hatjimihail AT. Relation of Diagnostic Accuracy Measures. Wolfram Demonstrations Project, Champaign: Wolfram Research, Inc., Champaign: Wolfram Research, Inc., 2018. Available at: <https://demonstrations.wolfram.com/RelationOfDiagnosticAccuracyMeasures/>

## License

[Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.](https://creativecommons.org/licenses/by-nc-sa/4.0/)

First Published: May 2, 2018